

OPTIMAL INSTRUMENT

Difficulty–Baseline

Instructions: Please circle the level of difficulty you have for each activity today.	Able to do without any difficulty	Able to do with little difficulty	Able to do with moderate difficulty	Able to do with much difficulty	Unable to do	Not applicable
1. Lying flat	1	2	3	4	5	9
2. Rolling over	1	2	3	4	5	9
3. Moving–lying to sitting	1	2	3	4	5	9
4. Sitting	1	2	3	4	5	9
5. Squatting	1	2	3	4	5	9
6. Bending/stooping	1	2	3	4	5	9
7. Balancing	1	2	3	4	5	9
8. Kneeling	1	2	3	4	5	9
9. Standing	1	2	3	4	5	9
10. Walking–short distance	1	2	3	4	5	9
11. Walking–long distance	1	2	3	4	5	9
12. Walking–outdoors	1	2	3	4	5	9
13. Climbing stairs	1	2	3	4	5	9
14. Hopping	1	2	3	4	5	9
15. Jumping	1	2	3	4	5	9
16. Running	1	2	3	4	5	9
17. Pushing	1	2	3	4	5	9
18. Pulling	1	2	3	4	5	9
19. Reaching	1	2	3	4	5	9
20. Grasping	1	2	3	4	5	9
21. Lifting	1	2	3	4	5	9
22. Carrying	1	2	3	4	5	9

23. From the above list, choose the 3 activities you would most like to be able to do without any difficulty (for example, if you would most like to be able to *climb stairs*, *kneel*, and *hop* without any difficulty, you would choose: 1. 13 2. 8 3. 14)

1. ____ 2. ____ 3. ____

24. From the above list of three activities, choose the primary activity you would most like to be able to do without any difficulty (for example, if you would most like to be able to *climb stairs* without any difficulty, you would choose: Primary goal, 13)

Primary goal. ____

© 2005, 2006 , 2012 American Physical Therapy Association. All rights reserved. No part of this instrument may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, or otherwise without prior permission of the American Physical Therapy Association. Contact permissions@apta.org or visit www.apta.org/publications.

Adapted/ revised in July 2005, August 2006, December 2012 with permission of APTA from Guccione AA, Mielenz TJ, De Vellis RF, et al. Development and testing of a self-report instrument to measure actions: Outpatient Physical Therapy Improvement in Movement Assessment Log (OPTIMAL). *Phys Ther.* 2005;85:515-530.

Confidence–Baseline

Instructions: Please circle the level of confidence you have for doing each activity today.	Fully confident in my ability to perform	Very confident	Moderate confidence	Some confidence	Not confident in my ability to perform	Not applicable
1. Lying flat	1	2	3	4	5	9
2. Rolling over	1	2	3	4	5	9
3. Moving–lying to sitting	1	2	3	4	5	9
4. Sitting	1	2	3	4	5	9
5. Squatting	1	2	3	4	5	9
6. Bending/stooping	1	2	3	4	5	9
7. Balancing	1	2	3	4	5	9
8. Kneeling	1	2	3	4	5	9
9. Standing	1	2	3	4	5	9
10. Walking–short distance	1	2	3	4	5	9
11. Walking–long distance	1	2	3	4	5	9
12. Walking–outdoors	1	2	3	4	5	9
13. Climbing stairs	1	2	3	4	5	9
14. Hopping	1	2	3	4	5	9
15. Jumping	1	2	3	4	5	9
16. Running	1	2	3	4	5	9
17. Pushing	1	2	3	4	5	9
18. Pulling	1	2	3	4	5	9
19. Reaching	1	2	3	4	5	9
20. Grasping	1	2	3	4	5	9
21. Lifting	1	2	3	4	5	9
22. Carrying	1	2	3	4	5	9

© 2005, 2006 , 2012 American Physical Therapy Association. All rights reserved. No part of this instrument may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, or otherwise without prior permission of the American Physical Therapy Association. Contact permissions@apta.org or visit www.apta.org/publications.

Adapted/ revised in July 2005, August 2006, December 2012 with permission of APTA from Guccione AA, Mielenz TJ, De Vellis RF, et al. Development and testing of a self-report instrument to measure actions: Outpatient Physical Therapy Improvement in Movement Assessment Log (OPTIMAL). *Phys Ther.* 2005;85:515-530.